

INTRAUTERINE BALLOON TAMPONADE-A NOVEL TECHNIQUE TO PREVENT AND MANAGE PLACENTAL SITE HAEMORRHAGE

Naila Tahir, Sadia Fatima, Muhammad Hamza Jahangeer*, Muhammad Adil, Shahbaz Khan

Combined Military Hospital Abbottabad Pakistan, *Banaras Hindu University Hafizabad Pakistan

ABSTRACT

Objective: This study was conducted to determine the effectiveness of Foley catheter balloon tamponade in postpartum haemorrhage at Combined Military Hospital (CMH) Abbottabad.

Study Design: Descriptive study.

Place and Duration of Study: Obstetrics & Gynaecology Department CMH Abbottabad, from Sep 2016 to Mar 2017.

Material and Methods: Patients who developed postpartum haemorrhage after caesarean section (CS) and were treated with intrauterine balloon tamponade (IUBT) using a Foley catheter were identified by review of medical records. Patient's demographic data, obstetric history, type of CS, cause of haemorrhage and transfusion history were extracted. Data was entered, coded and analyzed in IBM SPSS Statistics software. Descriptive statistics were produced.

Results: A total of 26 patients were included with mean age of 29.2 ± 5.23 years. More than two-thirds of the patients were multi-gravida and around three-quarters were at full term. The most frequent cause for postpartum haemorrhage (PPH) was placenta previa major. In 25 of the 26 (96.1%) patients the procedure was successful in stopping bleeding. There were no adverse effects related to the procedure.

Conclusion: We found that IUBT with an inexpensive Foley catheter is likely to be effective in most cases of PPH associated with caesarean section, without causing additional adverse events.

Keywords: Foley catheter, Intrauterine balloon tamponade, Postpartum haemorrhage.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Postpartum haemorrhage (PPH), defined as more than 500mL of blood loss within 24 hours of vaginal delivery or more than 1000mL after a caesarean section, is the leading direct cause of maternal mortality in the world¹. The World Health Organization (WHO) estimates that around a quarter of all maternal deaths are caused by PPH¹. PPH occurs in around 10% of all births but appropriate and timely management prevents death in most cases². The outcome, however, is worse in low-resource settings and PPH still has a significant negative impact in poor economies³. Around 31% of maternal deaths in Asia are due to PPH⁴ and it remains a significant health issue in Pakistan as well, where incidence is estimated to be 34%⁵. In comparison, the incidence of PPH in England is 13.8%⁶ and in the

U.S. it is estimated to be between 2% and 3%⁷.

Major risk factors for PPH include retained placenta or membranes, failure to progress during the second stage of labor, morbidly adherent placenta, lacerations, instrumental delivery, large for gestational age newborn, and hypertensive disorders⁷. For caesarean deliveries, general anesthesia, amnionitis, preeclampsia and prolonged active phase of labor have been reported to be significantly associated with PPH⁸. Most cases of PPH are primarily caused by uterine atony and the risk of death is dependent on general physical health of the mother especially the presence or absence of anemia. In addition, most deaths resulting from PPH occur during the first 24 hours after birth. Timely management is therefore extremely critical in terms of reducing mortality. Goals of treatment are to restore or maintain adequate circulatory volume to prevent hypoperfusion of vital organs, to restore or maintain adequate tissue

Correspondence: Dr Naila Tahir, Prof & HoD of Gynae Dept, CMH Abbottabad Pakistan (Email: dr-naila@hotmail.com)

Received: 20 Jul 2017; revised received: 08 Nov 2017; accepted: 22 Nov 2017

oxygenation, to reverse or prevent coagulopathy, and to eliminate the obstetric cause of PPH⁹.

Management options for PPH range from medical treatment and minimally invasive procedures to surgical interventions via laparotomy^{9,10}. Effective management of PPH usually requires a combination of interventions but options are limited for health facilities in low-resource settings. Intrauterine balloon tamponade (IUBT) has emerged as a promising option. If the balloon tamponade is effective in stopping the haemorrhage, the patient is less likely to require surgical intervention and blood transfusions, thereby decreasing the associated risks and cost¹⁰.

WHO has recognized IUBT as a significantly effective method for management of PPH,

catheter¹¹. Balloon catheters specifically designed for uterine tamponade are most widely used in developed countries. The high cost of these IUBT devices puts them out of reach for most low resource health systems or at least limits their use to a few selected hospitals in each country. Lower cost IUBT devices are therefore crucial for preventing mortality in these settings. This study was conducted to evaluate the efficacy of IUBT performed with the inexpensive 2-way Foley catheter in PPH occurring after caesarean delivery.

MATERIAL AND METHODS

This descriptive study was conducted as an internal evaluation at the department of Obstetrics & Gynecology of the Combined

Table-I: Obstetric characteristics of patients.

		Frequency	Percentage (%)
Gravidity	Primigravida	4	15.4
	Multigravida	18	69.2
	Grand multigravida	4	15.3
Parity	0	4	15.4
	1	3	11.5
	2	8	30.8
	3	7	26.9
	4	4	15.4
Term	Pre-term	7	26.9
	Full term	19	73.1
	Post term	0	0.0
Labor	None	23	88.5
	Failed induction	3	11.5

especially in low-resource areas, for both vaginal and caesarean deliveries². IUBT involves inserting a balloon device into the uterus and then incrementally filling the balloon with distilled water, which applies pressure to the uterus until the bleeding stops. Early use of IUBT can limit ongoing uterine blood loss and it is a simple procedure which can be readily implemented by providers with minimal training. Various types of IUBT devices are used including the Bakri Uterine Balloon, the Sengstaken-Blakemore Esophageal Tube, the Rusch Balloon, condom catheter and a Foley

Military Hospital (CMH) Abbottabad. The objective of the study was to compare our IUBT results with those reported in the literature and to share our results with other similar centres in the country. The study comprised of all patients who underwent a caesarean section between September 2016 and March 2017, developed PPH, and were treated with IUBT. The IUBT procedure involved using 2-way Foley catheter inflated with 80 to 100 mL of normal saline and retained for 24 hours. Non-probability convenience sampling was used. Total 26 patients were selected. The small sample size is due to

infrequent observation of uncontrolled PPH associated with caesarean section and therefore infrequent need for employing IUBT. Eligible patients were identified by review of medical charts of all patients admitted to the department during the study period and study data were extracted from the charts of eligible patients. Data extracted included patients' demographic data, obstetric history, type of caesarean section, ultrasono-graphic findings, cause of PPH, transfusions for PPH, and patient outcome. Data were entered, coded and analyzed in IBM SPSS Statistics software. Descriptive statistics were produced. Frequency and percentages were calculated for key study variables.

RESULTS

A total of 26 patients were included in the study who were between 21 and 36 years of

The most frequently observed ultrasonographic finding as well as the cause for PPH was placenta previa major. For all subjects with a normal situated placenta, uterine atony was determined to be the cause of PPH. Table-II presents the ultrasonographic findings and PPH cause.

Of the 26 patients treated, IUBT with 2-way Foley catheter inflated with 80 to 100 mL of normal saline and retained for up to 24 hours was successful in stopping PPH in 25 patients, a success rate of 96.2%. The patients in whom this intervention was not successful in cessation of haemorrhage, hysterectomy was performed. No mortality was observed among the patient group.

On average 2 (± 1.096) units of red blood cell concentrate (range 0 to 4) and 4.3 (± 2.290) units of fresh frozen plasma (range 0 to 9) were

Table-II: Ultrasonographic findings and cause of PPH.

		Frequency	Percentage (%)
Ultrasonographic Findings	Placenta previa major	14	53.8
	Placenta accreta	2	7.7
	Normal situated placenta	7	26.9
	Placenta previa major + increta	1	3.8
	Placenta previa major + accreta	2	7.7
Cause of PPH	Placenta previa major	14	53.8
	Placenta accreta	2	7.7
	Uterine Atony	7	26.9
	Placenta previa major + Increta	1	3.8
	Placenta previa major + accreta	2	7.7

age (mean age=29.2 years ± 5.23). Of the 26 caesarean sections, 19 (73.1%) were elective while 7 (26.9%) were emergency. Fourteen (53.8%) of the caesarean sections were performed under general anesthesia and 12 (46.2%) under spinal anesthesia. More than two-thirds of the patients were multi-gravida. Table-I presents the obstetric characteristics of the patients.

APGAR score after 1 minute for babies delivered via these caesarean sections ranged between 5 and 9, and the mean was 7.54 (± 0.86) whereas APGAR score after 5 minutes ranged between 8 and 10, mean being 9.54 (± 0.65).

transfused. Mean blood hemoglobin concentration before IUBT ranged from 8.5 to 13.6 mg/dL, mean being 11.1 mg/dL (± 1.39), whereas the mean blood hemoglobin concentration after treatment ranged from 7.6 to 12.6 mg/dL, mean being 10.1 mg/dL (± 1.32).

None of our patients suffered any adverse effects due to the IUBT procedure.

DISCUSSION

Due to the disadvantages like concealment of active bleeding and development of infection, uterine gauze packing is not preferred in the management of PPH. Uterine balloon temponade

was first reported by Goldrath in 1983¹². Since then balloon tamponade has emerged as a simple and effective intervention in PPH^{13,14}. Various different types of balloon catheters with various advantages and disadvantages have been employed for PPH. The Bakri postpartum balloon and the BT-cath balloon tamponade catheter are specifically designed for postpartum intrauterine tamponade however their cost could be prohibitive in low resource settings¹⁵. Other balloons can be used to achieve a similar effect in such settings. These include Sengstaken-Blakemore tube (used for treatment of bleeding esophageal varices), Rusch urologic balloon (used for stretching the bladder), and condom catheter (a condom is placed over the end of a Foley-type catheter, the base of the condom is tied to the catheter to prevent leakage, and then the condom is filled with up to 500mL fluid via the catheter)¹⁶. Rusch balloon and the condom catheter do not allow blood drainage from the uterine cavity. A 2-way Foley catheter has the advantage of being inexpensive as well as allowing blood drainage from the uterine cavity¹⁷. Efficacy of all these methods appears to be comparable. A comparison of IUBT using a condom-loaded Foley catheter with IUBT using Bakri balloon in management of PPH due to uterine atony following vaginal delivery found both of them equally effective in terms of success rates and maternal outcomes however Bakri balloon achieved homeostasis in a significantly shorter time¹⁸.

In a recent study¹⁹ on use of Foley catheter balloon tamponade to control placental site bleeding resulting from placenta previa major during caesarean section, of the 20 patients treated, PPH was successfully stopped in 17 (85%) patients while hysterectomy had to be performed for 3 (15%) patients. Our results are comparable to this study in terms of effectiveness.

A study done at Dow University of Health Sciences and Civil Hospital, Karachi evaluated the effectiveness of intrauterine balloon tamponade with a condom catheter in PPH due

to uterine atony following a vaginal delivery, instrumental delivery or caesarean section²⁰. Of the 139 patients treated, balloon tamponade was effective in stopping haemorrhage in 126 (90.4%) but was not effective in 13 (9.6%) patients. Our results of 96.2% effectiveness in cessation of PPH with a 2-way Foley catheter were comparable to the findings of this study.

The overall effectiveness rate reported in the literature for IUBT as a method of controlling PPH (regardless of the catheter type used) is between 80% and 100%²¹⁻²⁶. Our results are consistent with this reported overall rate of IUBT success.

CONCLUSION

We found that IUBT with an inexpensive Foley catheter is likely to be effective in most cases of PPH associated with caesarean section, without causing additional adverse events.

CONFLICT OF INTEREST

This study has no conflict of interest to declare by any author.

REFERENCES

1. Khan KS, Wojdyla D, Say L, Gülmezoglu AM, Van Look PFA. WHO analysis of causes of maternal death: a systematic review. *Lancet* 2006; 367(9516): 1066-74.
2. World Health Organization. WHO guidelines for the management of postpartum haemorrhage and retained placenta [Internet]. Geneva; 2009. Available from: http://www.who.int/reproductivehealth/publications/maternal_perinatal_health/9789241598514/en/
3. Ngwenya S. Postpartum Haemorrhage: Incidence, risk factors, and outcomes in a low-resource setting. *Int J Womens Health* 2016; 8: 647-50.
4. Gundabattula SR. Morbidly adherent placenta. *World clinics: Obstetrics & Gynecology: Postpartum haemorrhage* 2013; 2(2): 216-35.
5. Gani N, Ali TS. Prevalence and factors associated with maternal postpartum haemorrhage in Khyber Agency, Pakistan. *J Ayub Med Coll Abbottabad* 2013; 25(1-2): 81-5.
6. The Lancet. Countdown to 2030 for Reproductive, Maternal, Newborn, Child, and Adolescent Health and Nutrition [Internet]. 2015. Available from: <http://countdown2030.org/country-profiles/pakistan>
7. Hospital episode statistics: NHS Maternity Statistics - England, 2013-14 [Internet]. NHS Maternity Statistics - NHS Digital. 2015 [cited 2017]. Available from: <http://content.digital.nhs.uk/catalogue/PUB16725/nhs-mate-eng-2013-14-summ-repo-rep.pdf>
8. Suta J, Phaloprakarn C. A risk score for predicting postpartum haemorrhage in association with caesarean delivery. *Source J Obstet Gynaecol* 2015; 23(1): 3-11.

9. Belfont M. Overview of postpartum haemorrhage. In: Up To Date, Lockwood CJ (Ed), Up To Date, Waltham, MA. (Accessed on 2017).
10. Labarta FR, Recarte MP, Luque AA, Prieto LJ, Martín LP, Leyte MG, et al. Outcomes of pelvic arterial embolization in the management of postpartum haemorrhage: A case series study and systematic review. *Eur J Obstet Gynecol Reprod Biol* 2016; 206: 12-21.
11. Georgiou C. Balloon tamponade in the management of postpartum haemorrhage: a review. *BJOG* 2009; 116(6): 748-57.
12. Tirumuru S, Saba S, Morsi H, Muammar B. Intrauterine balloon tamponade in the management of severe postpartum haemorrhage: A case series from a busy UK district general hospital. *OJOG* 2013; 3: 131-6.
13. Sentilhes L, Vayssière C, Deneux-Tharoux C, Aya AG, Bayoumeu F, Bonnet MP, et al. Postpartum haemorrhage: Guidelines for clinical practice from the French College of Gynaecologists and Obstetricians (CNGOF): In collaboration with the French Society of Anesthesiology and Intensive Care (SFAR). *Eur J Obstet Gynecol Reprod Biol* 2016; 198: 12-21.
14. Sathe NA, Likis FE, Young JL, Morgans A, Carlson-Bremer D, Andrews J. Procedures and uterine-sparing surgeries for managing postpartum haemorrhage: A systematic review. *Obstet Gynecol Surv* 2016; 71(2): 99-113.
15. Tindell K, Garfinkel R, Abu-Haydar E, Ahn R, Burke TF, Conn K, et al. Uterine balloon tamponade for the treatment of postpartum haemorrhage in resource-poor settings: A systematic review. *BJOG* 2013; 120(1): 5-14.
16. Broekhuizen FF. A low-resource, simple method to treat postpartum haemorrhage. *Med Gen Med* 2003; 5(3): 39.
17. Seligman B, Xingzhu L. Economic assessment of interventions for reducing postpartum haemorrhage in developing countries. Cambridge, MA: Abt. Associates; 2006.
18. Darwish AM, Abdallah MM, Shaaban OM, Ali MK, Khalaf M, Sabra AM. Bakri balloon versus condom-loaded Foley's catheter for treatment of atonic postpartum haemorrhage secondary to vaginal delivery: A randomized controlled trial. *J Matern Fetal Neonatal Med* 2017; 31(6): 1-7.
19. Ali MK, Ahmed M, Abbas, Ahmed Y, Abdelbadee MS, Sherif A, et al. Use of Foley's catheter balloon tamponade to control placental site bleeding resulting from major placenta previa during caesarean section. *Proc Obstet Gynecol* 2016; 6(3): 4.
20. Lohano R, Haq G, Kazi S, Sheikh S. Intrauterine balloon tamponade for the control of postpartum haemorrhage. *J Pak Med Assoc* 2016; 66(1): 22-6.
21. Matsubara S, Takahashi H, Baba Y, Lefor AK. Re: Intrauterine balloon tamponade for management of severe postpartum haemorrhage in a perinatal network: A prospective cohort study. *BJOG* 2017; 124(11): 1792-93.
22. Martin E, Cheve MT, Legendre G, Multon O, Sentilhes L, Bouet PE. Maternal outcomes after uterine balloon tamponade for postpartum haemorrhage. *Obstetric Anesthesia Digest* 2016; 36(1): 32-3.
23. Kwon HY, Kim YH, Park YW, Kwon JY. Efficacy of bakri balloon tamponade in massive postpartum haemorrhage: A Series of 57 Cases. *Perinatology* 2016; 27(4): 236-43.
24. Tahaoglu AE, Balsak D, Erdogdu E, Bakır MS, Aksin S, Bala M, et al. Bakri balloon placement effectively treats uterine atony and placenta previa. *Biotechnol Biotechnol Equip* 2017; 31(4): 1-5.
25. Soyama H, Miyamoto M, Sasa H, Ishibashi H, Yoshida M, Nakatsuka M, et al. Effect of routine rapid insertion of Bakri balloon tamponade on reducing haemorrhage from placenta previa during and after caesarean section. *Arch Gynecol Obstet* 2017; 296(3): 469-74.
26. Kong MC, To WW. Balloon tamponade for postpartum haemorrhage: Case series and literature review. *Hong Kong Med J* 2013; 19(6): 484-90.