

FREQUENCY AND SEVERITY OF ANEMIA IN CHILDREN LESS THAN 15 YEARS OF AGE AT GWADAR DEVELOPMENT AUTHORITY HOSPITAL, GWADAR, BALUCHISTAN

Majid Latif, Saeed Bin Ayaz, Muneeba Manzoor*, Mansoor Ishaq

Combined Military Hospital Quetta/National University of Medical Sciences (NUMS) Pakistan, *Government Hospital Ghaziabad Lahore Pakistan

ABSTRACT

Objective: To determine the frequency and severity of anemia in children <15 years of age and find association of anemia with age and gender at Gwadar Development Authority Hospital, Gwadar, Baluchistan.

Study Design: Cross sectional study.

Place and Duration of Study: Gwadar Development Authority Hospital Gwadar, from Jun 2016 to Jun 2017.

Material and Methods: Through consecutive sampling, 483 children reporting to the pathology department for hemoglobin estimation were sampled. The samples were taken through antecubital veins using aseptic technique and the hemoglobin was analyzed using Medonic M-series M32 Hematology Analyzer. The sample was divided into three groups based on age i.e. group-1 (age 6-59 months), group-2 (age: 5-11 years), and group-3 (age: 11-14 years). We used definitions of anemia given by the World Health Organization. The data were analyzed using SPSS version 20.

Results: The mean age was 7.3 ± 4.3 years (range: 1-14 years). The frequency of anemia was 64.6% (n=312). Considering gender, the percentage of anemic children among male gender was 62.9% (n=132) and among females was 65.9% (n=180) ($p=0.483$). The percentage of anemic children was highest (74.5%) in group-1 ($p<0.001$). The gender could not find an association with anemia ($p=0.483$).

Conclusion: Anemia had a frequency of 64.6% in children under 15 years of age at Gwadar Development Authority Hospital, Gwadar. Age-group of 1-59 months had the highest percentage of anemic children. The gender did not have any association with anemia.

Keywords: Age, Anemia, Frequency, Gender, Gwadar, Hemoglobin, Severity.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Anemia is a condition in which the number of red blood cells are insufficient to meet physiologic needs of the body¹. Globally, anemia is a public health problem affecting people in both developed and developing countries with adverse effects on human health as well as on social and economic development². Anemia is a critical health concern as it adversely affects growth and energy levels³. According to the statistics of World Health Organization (WHO), worldwide, there are two billion people living with anemia⁴.

In developing countries, the estimated

prevalence of anemia is 39% in children less than 5 years of age and 48% in children 5-14 years of age⁴. In children, anemia affects motor and cognitive development and increases susceptibility to infections⁵. Lack of awareness among mothers, poor nutrition, unhealthy food habits, and parasitic infestations are the factors associated with lower hemoglobin levels in children⁶.

Although anemia remains a widespread public health problem in Pakistan, there are few studies on the prevalence and severity of anemia among children especially from Baluchistan. This was the first study carried out in Gwadar. The aim of this study was to describe the frequency and severity of anemia among pediatric patients under 15 years of age who underwent evaluation of their blood samples for hemoglobin levels in

Correspondence: Dr Majid Latif, Classified Pathologist, CMH Quetta Pakistan (Email: majidlatif33@hotmail.com)

Received: 06 Oct 2017; revised received: 29 Oct 2017; accepted: 27 Nov 2017

Gwadar development authority hospital, Gwadar. The association of anemia with the age and gender were secondary goals. The results of this study can be used by public health programmers to design target interventions aimed at reducing the huge burden of anemia in Gwadar.

MATERIAL AND METHODS

This was a cross-sectional study conducted at the pathology department, Gwadar development authority hospital gwadar, from June 2016 to June 2017 after taking permission from the hospital ethical committee. A sample size of 366 was estimated via Epi Tools Epidemiological calculator⁷ while keeping level of significance 5%, confidence level 95%, estimated true proportion 0.61%⁸, and 5% of absolute precision. Through consecutive sampling, we included 483 individuals who reported to the pathology department for hemoglobin estimation during the said time period.

After verbal informed consent, the samples were taken from the individuals through ante-cubital veins using aseptic technique. The samples were taken in potassium EDTA tube, uniformly mixed before analysis by blood tube mixer. The hemoglobin was analyzed using Medonic M-series M32 Hematology Analyzer (Boule Medical AB, Spanga, Sweden).

The sample was divided into three groups based on age i.e. group-1 (age 6-59 months), group-2 (age: 5-11 years), and group-3 (age: 11-14 years). We used definitions of anemia given by WHO given in table-I⁸.

The data were analyzed using SPSS version 20. The frequencies and percentages, means, and standard deviations were calculated for categorical and numerical data respectively. The mean hemoglobin levels among different genders and age-groups were compared using student t-tests and one-way analysis of variance with Games-Howell post hoc analysis. The association of anemia with gender and age-group was analyzed using Pearson's Chi-square analysis. A *p*-value <0.05 was considered significant.

RESULTS

The mean age was 7.3 ± 4.3 years with a median age of 8 years and a range of 1-14 years. Of 483 children finally included, 210 (43.5%) were male while 273 (56.5%) were female. There were 165 (34.2%) children in group-1 (age: 6-59 months), 201 (41.6%) children in group-2, and 117 (24.2%) children in group-3. The mean hemoglobin level in males was 10.6 ± 2.2 g/dl and in females was 10.6 ± 2 g/dl thus showing no significant difference ($p=1$). However, the eldest age group had the highest amount of hemoglobin ($p<0.001$). Using post hoc analysis, the mean hemoglobin level in group-3 was significantly greater than group-1 and group-2 ($p<0.001$ and $p=0.031$ respectively). The mean hemoglobin level in group -2 was significantly greater than that of group-2 ($p<0.001$) (table-II).

The frequency of anemia in the whole sample was 64.6% (n=312). Considering gender, the percentage of anemic children among male gender was 62.9% (n=132) and among females was 65.9% (n=180) ($p=0.483$). The percentage of anemic children was 74.5% (n=123) in group-1, 64.2% (n=129) in group-2, and 51.3% (n=60) in group-3 ($p<0.001$) (table-II).

DISCUSSION

Measurement of hemoglobin level is an important hematological parameter that helps to diagnose the extent and severity of anemia⁴. Numerous studies have demonstrated that even moderate anemia is associated with depressed mental and motor development in children and may not be reversible⁹. Pakistan has a developing menace of anemia, as most children, especially belonging to the low socioeconomic group are primarily anemic¹⁰. Numerous Pakistani studies have reported a prevalence of 61-78.7% in children¹⁰. The frequency observed by us (64.6%) also falls in the same range. Available data relating to the prevalence of anemia in Pakistan reveal that iron deficiency is the leading cause of all types of anemia in children¹⁰. The contributing factors of anemia have been identified as poverty, consumption of low iron diets, inappropriate

dietary habits, poor personal hygiene, and lack of sanitation¹¹.

The results of our study revealed that the frequency of anemia was highest in the age-group of 6-59 months. Our findings are consistent with the National Health survey for Pakistan conducted from 1990 to 1994, in which the prevalence of anemia was significantly associated with the child’s age, so that the youngest children had the highest odds of developing anemia¹². Similarly, a survey at Lahore had found highest

pregnancy and lactation. Anemic mothers usually give birth to children with moderate to severe anemia¹⁵. As children are exclusive breastfed during initial months, and the breast milk lacks adequate iron content, the increased prevalence of anemia in younger children is logical. At higher age, children start eating weaning foods such as meat, fish, or eggs¹⁵ that are rich in haem iron²², so the anemia improves. Secondly, since first two years of life are characterized by rapid growth demanding increased iron utilization, the risk of anemia particularly iron deficiency anemia

Table-I: Hemoglobin levels to diagnose anemia according to the World Health Organization guidelines.

Age-group	Normal hemoglobin levels in g/dL	Anemia		
		Mild	Moderate	Severe
		Hemoglobin in g/dL		
Children 6-59 months of age	11.0 or higher	10.0-10.9	7.0-9.9	lower than 7.0
Children 5-11 years of age	11.5 or higher	11.0-11.4	8.0-10.9	lower than 8.0
Children 11-14 years of age	12.0 or higher	11.0-11.9	8.0-10.9	lower than 8.0

Table-II: Frequency and severity of anemia in different age groups.

	Groups based on age		
	Group 1 (6-59 months)	Group 2 (5-11 years)	Group 3 (11-14 years)
Mean hemoglobin levels in g/dl	Mean ± standard deviation		
	9.7 ± 1.9	10.7 ± 1.7	11.3 ± 2.3
Sample groups based on hemoglobin levels	n (%)		
Normal value	42 (25.5)	72 (35.8)	57 (48.7)
Mild Anemia	39 (23.6)	33 (16.4)	27 (23.1)
Moderate anemia	66 (40)	78 (38.8)	21 (17.9)
Severe anemia	18 (10.9)	18 (9)	12 (10.3)

percentage of anemic children in age group of 1-4 years¹³. Contrary to this, a Pakistani study from Khyber Pakhtunkhwa has found highest percentage of anemia in the eldest age-group of 10-12 years among the sampled children¹⁴.

In India, highest prevalence of anemia has been observed in children below 5 years of age¹⁵. Studies from Brazil, Nepal, Bangladesh, Iran, India, Philippines, and Ethiopia have also endorsed younger age as an important risk factor associated with anemia¹⁶⁻²¹. The reason for increased prevalence in younger children can be related to the higher incidence of anemia during

is increased.

We did not find association of anemia with gender of the sampled children. Similar findings were observed by *Kishwar et al* in a study conducted at Lahore²³. However, other studies for Pakistan have found an association of anemia with gender with some reporting higher incidence in female children^{14,24} and others, a higher incidence in male children^{12,13}. There are conflicting evidences in the international data with regard to the relationship between gender and anemia in children. Studies from Yemen and India found a higher prevalence of anemia in

girls than boys^{25,26} and studies from Kenya and Haiti, and some from India found boys to be more at risk²⁷⁻²⁹.

The data in our study and other studies conducted in Pakistan may provide valuable insight into the prevalence of anemia in children. Considering the burden of anemia in Pakistani children, it is essential that interventions such as iron supplementation and food fortification should be used at mass scale. Currently wheat flour fortification and iron supplementation are successful strategies being used worldwide³⁰. Furthermore, recent studies in Central Asia, Venezuela, and Iran suggest that wheat flour fortification can significantly improve iron status at the population level³¹ but there is a lack of such supplementation programs in Pakistan. High frequency of anemia in children needs due emphasis so as to bring down the total prevalence of anemia. Among school-age children and adolescent boys, those who have a history of iron deficiency anemia, special health-care needs, or low iron intake should periodically be screened for anemia and appropriate measures be taken³⁰.

It was a cross-sectional study so the temporal relationship between anemia and the associated factors could not be established. Moreover, the study lacked detailed investigation of the morphological appearance of red blood cells to differentiate various types of anemia.

CONCLUSION

The frequency of anemia in the sampled children under 15 years of age in Gwadar Development Authority Hospital, Gwadar was 64.6%. Age-group of 1-59 months had the highest percentage of anemic children. The gender did not have any association with anemia.

CONFLICT OF INTEREST

This study has no conflict of interest to declare by any author.

REFERENCES

1. Assessing the iron status of populations: Report of a joint World Health Organization/Centers for disease control and prevention technical consultation on the assessment of iron status at the

- population level, 2nd ed, Geneva, World Health Organization, 2007. Available at: http://www.who.int/nutrition/publications/micronutrients/anaemia_iron_deficiency/9789241596107.pdf
2. World Health Organization. The World Health Report 2002: Reducing risks, promoting healthy life. Geneva: World Health Organization; 2002: 49-93.
3. United Nations Children's Fund/United Nations University/WHO: Iron deficiency anaemia. Assessment, prevention and control. A guide for programme managers (WHO/NHD/01.3). Geneva: World Health Organization 2001; 15-31.
4. WHO, UNICEF, UNU, Iron deficiency anaemia: Assessment, prevention and control, a guide for programme managers. Geneva, World Health Organization, 2001. Available from: http://www.who.int/nutrition/publications/micronutrients/anaemia_iron_deficiency/WHO_NHD_01.3/en/index.html
5. Baker RD, Greer FR. Committee on Nutrition American Academy of Pediatrics. Diagnosis and prevention of iron deficiency and iron-deficiency anemia in infants and young children (0-3 years of age). *Pediatrics* 2010; 126(5): 1040-50.
6. Assefa S, Mossie A, Hamza L. Prevalence and severity of anemia among school children in Jimma Town, Southwest Ethiopia. *BMC Hematol* 2014; 14(1): 3.
7. Epi Tools Epidemiological Calculators. AusVet. Accessed 2016. Available from: <http://epitools.ausvet.com.au/content.php?page=1Proportion>.
8. Stevens GA, Finucane MM, De-Regil LM, Paciorek CJ, Flaxman SR, Branca F, et al. Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995-2011: A systematic analysis of population-representative data. *Lancet Glob Health* 2013; 1(1): e16-25.
9. Grantham-McGregor S, Ani C. A review of studies on the effect of iron deficiency on cognitive development in children. *J Nutr* 2001; 131(2S-2): 649S-68S.
10. Akhtar S, Ahmed A, Ahmad A, Ali Z, Riaz M, Ismail T. Iron status of the Pakistani population-current issues and strategies. *Asia Pac J Clin Nutr* 2013; 22(3): 340-7.
11. Agha F, Sadaruddin A, Khan RA, Ghafoor A. Iron deficiency in adolescents. *J Pak Med Assoc* 1992; 42(1): 3-5.
12. Habib MA, Black K, Soofi SB, Hussain I, Bhatti Z, Bhutta ZA, et al. Prevalence and predictors of iron deficiency anemia in children under five years of age in Pakistan, A secondary analysis of national nutrition survey data 2011-2012. *PLoS One* 2016; 11(5): e0155051.
13. Nadeem R, Ahmed N, Bashir S. Frequency of iron deficiency anaemia and beta thalassaemia trait at haematology department of children hospital, Lahore. *Biomedica* 2012; 28(1): 66-70.
14. Ullah I, Zahid M, Sthanadar AA, Sthanadar IA, Ali PA, Shah M, et al. Iron deficiency anemia in school age children in district Karak, Khyber Pakhtunkhwa province, Pakistan. *Open J Blood Dis* 2014; 4(2): 9-15.
15. Alvarez-Uria G, Naik PK, Midde M, Yalla PS, Pakam R. Prevalence and severity of anaemia stratified by age and gender in rural India. *Anemia* 2014; 2014: 176182.
16. Ghosh A, Ghartimagar D, Thapa S, Sathian B, De A. Microcytic Hypochromic Anemia in Pediatric Age Group: A Hospital Based Study in Nepal. *Am J Pub Health Res* 2015; 3(4A): 57-61.
17. Zuffo CR, Osório MM, Taconeli CA, Schmidt ST, da Silva BH, Almeida CC. Prevalence and risk factors of anemia in children. *J Pediatr Rio J* 2016; 92(4): 353-60.
18. Uddin MK, Sardar MH, Hossain MZ, Alam MM, Bhuya MF, Uddin MM, et al. Prevalence of anaemia in children of 6 months to 59 months in Narayanganj, Bangladesh. *J Dhaka Med Coll* 2010; 19(2): 126-30.

19. Keikhaei B, Zandian K, Ghasemi A, Tabibi R. Iron-deficiency anemia among children in southwest Iran. *Food Nutr Bull* 2007; 28(4): 406-11.
 20. Tengco LW, Rayco-Solon P, Solon JA, Sarol JN, Solon FS. Determinants of anemia among preschool children in the Philippines. *J Am Coll Nutr* 2008; 27(2): 229-43.
 21. Gebreegziabihier G, Etana B, Niggusie D. Determinants of Anemia among children aged 6-59 months living in Kilte Awulaelo Woreda, Northern Ethiopia. *Anemia* 2014; 2014: 245870.
 22. Abbaspour N, Hurrell R, Kelishadi R. Review on iron and its importance for human health. *J Res Med Sci* 2014; 19(2): 164-74.
 23. Kishwar F, Ashraf T, Hanif I, Hanif A, Kalsoom S. Iron deficiency anemia; prevalence in children of Lahore. *Prof Med J* 2015; 22(9): 1122-5.
 24. Sheikh S, Shaikh KR, Memon M. Prevalence of iron deficiency and thalassemia trait in children at pediatric department of L.U.H. *Biomedica* 2003; 19: 11-7.
 25. Al-Zabedi EM, Kaid FA, Sady H, Al-Adhroey AH, Amran AA, Al-Maktari MT. Prevalence and risk factors of iron deficiency anemia among children in Yemen. *Am J Health Res* 2014; 2(5): 319-26.
 26. Kaur IP, Kaur S. A comparison of nutritional profile and prevalence of anemia among rural girls and boys. *J Exerc Sci Physio Ther* 2011; 7(1): 11-8.
 27. Foote EM, Sullivan KM, Ruth LJ, Oremo J, Sadumah I, Williams TN, et al. Determinants of anemia among preschool children in rural, western Kenya. *Am J Trop Med Hyg* 2013; 88(4): 757-64.
 28. Ayoya MA, Ngnie-Teta I, Séraphin MN, Mamadoultabou A, Boldon E. Prevalence and risk factors of anemia among children 6-59 months old in haiti. *Anemia* 2013; 2013: 502968.
 29. Pasricha SR, Black J, Muthayya S, Shet A, Bhat V, Nagaraj S, et al. Determinants of anemia among young children in rural India. *Pediatrics* 2010; 126(1): e140-9.
 30. Pasricha SR, Drakesmith H, Black J, Hipgrave D, Biggs BA. Control of iron deficiency anemia in low- and middle-income countries. *Blood* 2013; 121(14): 2607-17.
 31. Sadighi J, Mohammad K, Sheikholeslam R, Amirkhani MA, Torabi P, Salehi F, et al. Anaemia control: lessons from the flour fortification programme. *Public Health* 2009; 123(12): 794-9.
-