

RISK FACTORS ASSOCIATED WITH IN-HOSPITAL DEATH IN ADULT COVID-19 PATIENTS IN KARACHI, PAKISTAN: A RETROSPECTIVE CHART REVIEW

Imran Ahmad, Farooq Azam Rathore, Muhammad Nadir Khan, Shaikh Nabi Bukhsh Nazir*

Pakistan Naval Ship (PNS) Shifa Hospital, Karachi Pakistan, *Dow University of Health Sciences, Karachi Pakistan

ABSTRACT

Objective: To describe an audit of COVID-19 related mortality in the largest Navy hospitals in Pakistan and identify the risk factors associated with in-hospital death in these patients.

Study Design: Retrospective chart review.

Place and Duration of Study: Department of Medicine, Pakistan Naval Ship (PNS) Shifa Hospital, Karachi, from 1st Apr to 30th May 2020.

Methodology: All patients who died in the Intensive care unit (ICU) of PNS Shifa Hospital, Karachi during study period, and having PCR positive COVID-19 infection were included. The data recorded included basic demographics (age, gender and length of stay in hospital), documentation of co-morbidities (e.g. diabetes mellitus [DM], ischemic heart disease [IHD] and other medical and surgical problems), need for mechanical ventilation and most probably cause or contributory factor to the death of the patient. Descriptive statistics were used for calculating mean and frequencies. Chi-square test was used to determine the association between variables of the study.

Results: A total of 22 cases were included. There were 21 males. The mean age was 65.7 years. Most 12 (56.5%) of the patients were more than 65 years of age. DM was the most common risk factor in 15 (69%) patients followed by hypertension 13 (59%) and IHD in 10 (45%). The hospital admission ranged from 1 to 19 days with a mean of 7.13 days.

Conclusion: We identified length of stay, mechanical ventilation, age >65 years, pneumonia and DIC are the potential risk factors for in-hospital mortality in severely ill COVID-19 patients. This audit can guide the clinicians working in resource constrained areas, in making decision regarding hospital admission for the high-risk cases.

Keywords: Coronavirus, Complication, Death, Intensive care, Mortality.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

The novel coronavirus (COVID-19) originated in December 2019 from Hunan sea food market, China and rapidly spread across the globe in the form of a pandemic affecting more than 200 countries¹. The World Health Organization (WHO) has reported around 9.4 million cases all around the globe with a death toll more than 484,000². The number of deaths all over the world is unprecedented considering the medical advances of the 21st century.

COVID-19 primarily affects the respiratory system but other organs including cardiovas-

cular, renal, gastrointestinal tract and central nervous system may also be involved^{3,4}. Predominant cause of death in COVID-19 documented in the literature is respiratory failure, which may be further complicated by sepsis, Acute respiratory distress syndrome (ARDS), and micro thrombi in pulmonary vasculature. However, death from acute myocardial infarctions, myocarditis, sudden death, strokes, and cytokine release syndromes leading to kidney injury, disseminated intravascular coagulation (DIC) and multi organ dysfunction have also been reported⁵⁻⁷.

The case fatality of COVID-19 has been reported to range from 1% to 7% depending upon the population screened. In countries like South Korea and Switzerland, where mass screening was done it was documented as 1%, since mild

Correspondence: Dr Farooq Azam Rathore, Assistant Professor, Dept of Rehabilitation, PNS Shifa Hospital, Karachi Pakistan
Email: farooqrathore@gmail.com

Received: 29 Jun 2020; revised received: 15 Jul 2020; accepted: 27 Jul 2020

or asymptomatic cases were also included in the data reporting. However, in countries like Spain and Italy where limited population screening was performed or only people presenting to hospital were screened it was reported as 7%⁸. The official number of confirmed cases in Pakistan is around 198,800 with 4000 deaths (27th June 2020 estimates)⁹. There has been an unprecedented number of publications on different aspects of COVID-19 mostly from China, Europe and the USA. There is a paucity of data from the developing countries including Pakistan which have seen a rapid rise in the number of cases in the last 4 weeks.

The aim of the paper is to provide a mortality audit of COVID-19 related death in one of the largest Navy hospitals in Pakistan and to identify the risk factors for in-hospital mortality in this population of patients. This is part of an ongoing prospective study, and we are reporting the experience of first 08 weeks.

METHODOLOGY

A retrospective chart review of patient's admitted to the Intensive care unit (ICU) of PNS Shifa Hospital, Karachi between 1st April to 30th May 2020, for COVID-19 infection was conducted. The study was approved by the hospital ethics review committee (ERC/2020/REHAB/26, June 2020). A dedicated COVID-19 clinic has been established for the screening of suspected cases and residents are available in the emergency department round the clock to check, admit and treat suspected or confirmed cases of COVID-19.

Only seriously ill patients who are unable to maintain oxygen saturation or are rapidly deteriorating are admitted in the ICU. The patient record at the PNS Shifa is maintained manually in paper files. The data collection was performed by the lead author (IA) and was extracted on the pre-designed data extraction sheet. We only included confirmed cases of COVID-19 who expired in the ICU till 30th May 2020. A sample of 22 patients was collected using the convenience sampling technique. Diagnosis was made on Polymerase Chain Reaction (PCR) of the nasopharyngeal swab sample. The national and inter-

national guidelines for management of severely ill COVID-19 patients were followed in all cases. The data recorded included basic demographics (age, gender and length of stay in hospital), documentation of co-morbidities (e.g. diabetes mellitus, ischemic heart disease and other medical and surgical problems), need for mechanical ventilation and most probably cause or contributory factor to the death of the patient. Data was computed and analyzed using SPSS 23.0 (IBM, Armonk, NY, USA). Percentages and frequencies were calculated to present a summary of baseline study parameters and Mean \pm SD were presented for quantitative variables. The chi-square test was used to investigate the level of association among variables. A $p \leq 0.05$ was set as threshold to detect statistical significance.

RESULTS

There were 25 mortalities in the ICU during this time. We identified 22 confirmed cases of COVID-19 who expired in the ICU of PNS Shifa during the study period. Three cases were excluded after a negative PCR test and twenty-two cases were included in the final review (table-I). There were 21 males and 1 female. The mean age was 65.7 ± 10.25 years with an age range of 50-84 years. Half (12/22) of the patients were above 65 years of age. Diabetes mellitus (DM) was the most common risk factor documented in 15/22 (69%) patients followed by hypertension (HTN) in 13 (59%) and ischemic heart disease (IHD) in 10 (45%). One patient had pre-existing chronic liver disease, and another was a post-operative case who developed COVID-19 related complications during hospital admission and ultimately expired in the ICU. More than one risk factors (any combination of DM, IHD and HTN) was documented in 16 (72%) patients and eight patients had three risk factors (table-I).

The cause of death in COVID-19 was mostly MODS 10 (45.45%), followed with ARDS 8 (36.36%), sudden cardiac death 7 (31.81%), respiratory failure 6 (27.27%) and myocarditis 3 (13.63%). Table-II presents the association of patient characteristics with COVID-19 cause of

death. 8/22 patients died while on ventilator support 10/22 patients did not consented for ventilator support to 19 days with a mean of 7.13 days. Statistically significant difference was noted in age classes

Table-I: Clinical features and demographics of patients presented in this case series.

S. No.	Gender	Age (Yrs)	Duration of Admission (Days)	Co-morbidities and associated issues	Mechanical Ventilation	Complications	Cause of Death
1	Female	68	4	DM, HTN, IHD	Family did not consent for Mechanical Ventilation	Progressive respiratory failure	ARDS/ Respiratory failure DNR
2	Male	69	13	DM, HTN	Initially placed on Mechanical Ventilation but then shifted to NIV	Myocarditis with ejection fraction of 40%, DIC, Cytokine storm	Myocarditis and MODS
3	Male	60	5	DM, HTN, IHD, CKD	Yes	Myocarditis, Acute on chronic Renal failure	ARDS
4	Male	66	1	Liver Cirrhosis	No	Nil	Sudden cardiac arrest
5	Male	84	11	Obstructed hernia, HTN	Family did not consent for Mechanical Ventilation	Severe COVID-19 pneumonia with MODS	MODS/DNR
6	Male	76	01	DM, HTN, IHD Parkinson disease	Family did not consent for Mechanical Ventilation	Respiratory failure and MODS	MODS/DNR
7	Male	80	02	COPD, Bilateral subdural hematoma	Family did not consent for Mechanical Ventilation	Progressive respiratory failure	Progressive respiratory failure/DNR
8	Male	73	01	DM, HTN, IHD	Family did not consent for Mechanical Ventilation	Progressive respiratory failure	Sudden cardiac arrest after admission
9	Male	52	10	HTN, IHD, CVA	Family did not consent for Mechanical Ventilation	COVID-19 pneumonia with ARDS	Sudden cardiac arrest
10	Male	66	19	DM, HTN	Yes	Severe COVID-19 pneumonia, ARDS, Ventilator associated pneumonia Sepsis and sudden cardiac arrest.	MODS
11	Male	60	3	DM, HTN	Yes	Severe COVID-19 pneumonia, Ventilator associate pneumonia ARDS	MODS
12	Male	55	7	DM, HTN	Yes	Cytokine storm syndrome, Severe COVID-19 pneumonia, ARDS, Acute kidney injury and septic shock	MODS
13	Male	55	4	Nil	Yes	Severe pneumonia and ARDS	MODS
14	Male	58	13	DM	Yes	Severe COVID-19 pneumonia, ARDS, Septic shock Cytokine storm syndrome	MODS
15	Male	55	9	DM, HTN, IHD	Yes	Severe COVID-19 pneumonia ARDS Septic shock Cytokine storm syndrome	MODS
16	Male	72	01	DM, HTN, IHD	Individual did not consent for mec-hanical ventilation	Severe COVID-19 pneumonia	Progressive respiratory failure
17	Male	50	09	HTN, IHD	Yes	Severe COVID-19 pneumonia ARDS Myocarditis hypotensive shock DIC Acute kidney injury	MODS
18	Male	70	04	DM, HTN	Individual did not consent for mecha-nical ventilation	Severe COVID-19 pneumonia	Progressive Respiratory failure
19	Male	82	13	DM, IHD	Initially placed on vent and weaned off	Sudden hypotensive shock followed by cardiac arrest	Sudden cardiac arrest
20	Male	61	10	COPD Fecal peritonitis due to intestinal perforation with Fournier gangrene	Individual did not consent for mechanical ventilation	Fecal peritonitis leading to sepsis and cardiac arrest	Sudden cardiac arrest
21	Male	78	13	DM, IHD	Individual did not consent for mechanical ventilation	Severe COVID-19 pneumonia with MODS	MODS
22	Male	56	04	DM	No		Sudden cardiac arrest

ventilator support and were either on NIV or with ARDS and respiratory failure. Ventilatory oxygen. The hospital admission ranged from 1 status had also significantly association with

MODS, sudden cardiac death and respiratory failure.

Table-III shows the association of deaths noted in COVID-19 with co-morbidities and complications occurred during the plan of care in hospital. Pneumonia was significantly associated with causes of death including ARDS and MODS

in COVID-19. DIC was also significantly associated with ARDS. Postmortem examination was not carried out in any case.

DISCUSSION

Management of COVID-19 is a global health challenge both for the high and low income countries. Despite rapid sharing of information,

Table-II: Association between patient's characteristics and cause of death in COVID-19.

		ARDS	Respiratory Failure	Myocarditis	MODS	Sudden Cardiac Death
Age	50-60	6 (75%)	-	3 (100%)	6 (60%)	2 (28.6%)
	61-71	2 (25%)	2 (33.3%)	-	1 (10%)	3 (42.9%)
	>71	-	4 (66.7%)	-	3 (30%)	2 (28.6%)
	<i>p</i> -value	0.022*	0.036*	0.081	0.161	0.514
Gender	Male	7 (87.5%)	5 (83.3%)	3 (100%)	10 (100%)	7 (100%)
	Female	1 (12.5%)	1 (16.7%)	-	-	-
	<i>p</i> -value	0.364	0.273	>0.9	>0.9	>0.9
Length of stay	1-6	2 (25%)	6 (100%)	2 (66.7%)	4 (40%)	3 (42.9%)
	7-12	4 (50%)	-	1 (33.3%)	3 (30%)	2 (28.6%)
	13-18	1 (12.5%)	-	-	2 (20%)	1 (14.3%)
	>18	1 (12.5%)	-	-	1 (10%)	1 (14.3%)
	<i>p</i> -value	0.115	0.041*	0.794	0.648	0.505
Mechanical Ventilation	Family did consent	2 (25%)	6 (100%)	-	3 (30%)	3 (42.9%)
	Yes	6 (75%)	-	2 (66.7%)	7 (70%)	2 (28.6%)
	No	-	-	1 (33.3%)	-	2 (28.6%)
	<i>p</i> -value	0.095	0.007*	0.124	0.08*	0.08*

Table-III: Association between patients' complication, co-morbidity and cause of death in COVID-19.

	ARDS	Respiratory Failure	Myocarditis	MODS	Sudden Cardiac Death
DM	5 (62.5%)	5 (83.3%)	2 (66.7%)	8 (80%)	4 (57.1%)
<i>p</i> -value	>0.9	0.616	>0.9	0.630	0.387
HTN	6 (75%)	5 (83.3%)	2 (66.7%)	7 (70%)	4 (57.1%)
<i>p</i> -value	>0.9	0.616	>0.9	>0.9	0.630
IHD	4 (50%)	4 (66.7%)	2 (66.7%)	4 (40%)	3 (42.9%)
<i>p</i> -value	>0.9	0.348	0.571	0.691	>0.9
CKD	-	-	1 (33.3%)	1 (10%)	-
<i>p</i> -value	-	-	0.136	0.455	-
Liver Cirrhosis	-	-	-	-	1 (14.3%)
<i>p</i> -value	-	-	-	-	0.317
Stroke	1 (12.5%)	-	-	-	1 (14.3%)
<i>p</i> -value	0.364	-	-	-	>0.9
COPD	-	1 (16.7%)	-	-	1 (14.3%)
<i>p</i> -value	-	0.481	-	-	>0.9
Pneumonia	8 (100%)	4 (66.7%)	1 (33.3%)	9 (90%)	2 (28.6%)
<i>p</i> -value	0.018*	>0.9	0.527	0.031*	0.052
DIC	4 (50%)	-	1 (33.3%)	3 (30%)	-
<i>p</i> -value	0.039*	0.266	>0.9	0.624	-

there are still many unanswered questions and lack of clear evidence for many treatments. Although lot of information is now available on the clinical presentation and basic epidemiology of COVID-19, there is still a paucity of COVID-19 mortality data especially from the developing countries. The overall case fatality rate for COVID-19 is reported to be 4.5% but ICU related mortality is remarkably high ranging from 30-70%. Remarkably high mortality rates have

results in a cytokine storm syndrome and MODS leading to death. During the cytokine storm and oxidative stress, a hypercoagulable state occurs in the body with increased risk of venous and arterial embolism. This may be further complicated by development of DIC which is associated with 27 times increased chances of death. A gender difference in COVID-19 related mortality has been reported and males are more severely affected. This is likely related to the higher exp-

Table-IV: Comparison of this series with selected COVID-19 mortality reports from Wuhan, China and Italy.

Parameters	Zhou <i>et al</i>	Du <i>et al</i>	Onder <i>et al</i>	Current Series
Number of cases	54	21	355	22
Country	China	China	Italy	Pakistan
World Bank Income Based Classification	Upper-Middle Income Country	Upper-Middle Income Country	High Income Country	Low-Middle Income Country
Centers	Jinyintan Hospital and Wuhan Pulmonary Hospital (Wuhan, China)	Wuhan Pulmonary Hospital (Wuhan, China)	Country level data	PNS Shifa Hospital (Karachi)
Median age (Years)	69	70	79.5	65.72
Gender	38 (70%) M	10 (47.6%) F	106 (30.0%) F	21 (95.5%) M
Hypertension	26 (48%)	13 (61.9%)	Nil	13 (59%)
Diabetes mellitus	17 (31%)	6 (28.6%)	126(35.5%)	15 (68.2%)
Ischemic heart disease	13 (24%)	12 (57.1%)	117 (30%)	10 (45.5%)
Chronic obstructive lung disease	4 (7%)	-	-	2 (9%)
Chronic kidney disease	2 (4%)	1 (4.7%)	-	01 (4.5%)
Chronic liver disease	-	1 (4.7%)	-	01 (4.5%)
Post-operative case	-	-	-	01 (4.5%)
Active cancer	-	-	72 (20.3%)	-
Stroke	-	-	34 (9.6%)	1 (4.7%)

been documented in the elderly and patients with comorbidities¹⁰⁻¹². Therefore, it becomes imperative to have local data to guide policies and device management strategies to identify high risk patients.

SARS-COV-2 attaches to the host cell through the Angiotensin-converting enzyme 2 (ACE-2) receptor leading to internalization and subsequent replication of the virus. During that phase a dysregulated immune response is generated which leads to hyper inflammation and release of mediators of inflammation including complement and Interleukin-6 (IL-6)¹³. This

ression of ACE-2 receptor in males as compared to females and the lack of the protective effect of estrogen and X chromosome¹⁴.

Zhao *et al* performed a retrospective cohort study of COVID-19 related mortality at 2 different hospitals in Wuhan, China¹⁵. There are some similarities and differences with our cohort (table-IV). The mean age in our cohort is higher than the mean age reported by Zhao *et al* (65 years vs. 54.5 years) Males constitutes the pre-dominant group however we had very few females in our cohort than their group. This could be attributed to less severity of COVID-19

in females, but such low numbers could also represent the attitude of society towards females where males are given priority than females. They reported HTN (19%) as the most common risk factor followed by DM (8.2%), and cardiovascular diseases (CVD) in 2.7%. Whereas we documented diabetes mellitus as the most common risk factor in 15 (69%), followed by HTN in 13 (59%) and IHD in 10 patients. We are not sure about the high prevalence of diabetes in our cohort. One explanation can be that due to low level of awareness in the general public, combined with low literacy rates and inadequate follow-up by patients the incidence of uncontrolled DM is higher in Pakistan and might have contributed to a higher number in this study¹⁶.

Our data has certain similarities and differences with the data reported from other countries (table-IV). Zhao and Colleagues predicted that age more than 60 years, male gender and any comorbid including DM, HTN and cerebrovascular disease are risk factors for increased mortality in COVID-19 patients. We also observed higher mortality in males along with an increased prevalence of DM and HTN in our cohort.

Du *et al* presented the mortality audit of 22 patients from Wuhan, China. They reported HTN in 13 (61.9%), DM in 6 (28.6%) and IHD in 12 (57.1%) patients which is quite similar to our findings¹⁷.

Onder and colleagues studied the predictors of mortality in a nationwide Italian cohort of 355 patients¹⁸. The mean age in their report was 79 years as compared to 65.7 years in our cohort. One possible explanation is that the life expectancy of 83 years in Italy is much better than 67.7 years for Pakistan. However, one important finding is the presence of risk factors in large number of their cohort. Eighty-nine (25.1%) had a one disease, 91 (25.6%) had 2 diseases, and 172 (48.5%) had 3 or more associated diseases. In comparison, patients in our study had two risk factors in 16 (72%) and had three risk factors in 8 (36%).

Mortality in our case series was predominantly due to MODS in ten cases. Six patients expired due to sudden cardiac death although we cannot comment if it was due to arrhythmias or pulmonary embolism. However, this could represent myocarditis or heart failure which is being identified as an important cause of mortality in COVID-19. Five deaths in our study were due to ARDS. Ruan and colleagues from Wuhan, China reported respiratory failure in 53% of their cohort followed by myocardial damage in 7% and both in 33% cases¹⁹. The mean duration of admission in our cases was 7 days and 14/22 patients died within ten days.

LIMITATION OF STUDY

Considering the global mortality of COVID-19, this is a relatively small case series. But it adds useful data to the global pool of knowledge being the first attempt of COVID-19 mortality audit from Pakistan. It was a retrospective review and included only risk factors for in-hospital mortality. There is a need for detailed documentation and reporting of the laboratory and radiological parameters in future COVID-19 mortality audits. In addition, this chart review did not document the possible treatment related complications which can be critical in-patient evaluation and outcome.

CONCLUSION

This single center mortality audit suggests that increasing age with the presence of pneumonia, length of stay and DIC are at an increased risk of death. There is also an increased the risk of death for patients who develop ARDS, cardiac injury and acute kidney disease. We hope that this experience of a major tertiary care hospital for COVID-19 related mortality will guide the clinicians in making decision regarding hospital admission for the high-risk cases. This will likely identify cases at higher risk for mortality so that an early and aggressive workup and treatment plan can be instituted, and resources can be prioritized during this global health care crisis. This is especially important for resource constrained areas like Pakistan.

CONFLICT OF INTEREST

The authors declared no conflict of interest related to the study.

REFERENCES

1. Ge H, Wang X, Yuan X, Xiao G, Wang C, Deng T, Yuan Q, Xiao X. The epidemiology and clinical information about COVID-19. *Eur J Clin Microbiol Infect Dis* 2020; 39(1): 1011-19.
2. World Health Organization. Coronavirus disease (COVID-19). Situation Report-158. Available at https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200626-covid-19-sitrep-158.pdf?sfvrsn=1d1aae8a_2
3. Ahmad I, Rathore FA. Neurological manifestations and complications of COVID-19: A literature review. *J Clin Neurosci* 2020; 77(1): 8-12.
4. Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020; 395(10223): 497-506.
5. Vincent JL, Taccone FS. Understanding pathways to death in patients with COVID-19. *Lancet Respir Med* 2020; 8(1): 430-32.
6. Li X, Xu S, Yu M, Wang K, Tao Y, Zhou Y, et al. Risk factors for severity and mortality in adult COVID-19 inpatients in Wuhan. *J Allergy Clin Immunol* 2020; S0091(20): 30495-4.
7. Wang K, Zhang Z, Yu M, Tao Y, Xie M. 15-day mortality and associated risk factors for hospitalized patients with COVID-19 in Wuhan, China: an ambispective observational cohort study. *Intensive Care Med* 2020; 46(7): 1-3.
8. Martins-Filho PR, Tavares CSS, Santos VS. Factors associated with mortality in patients with COVID-19. A quantitative evidence synthesis of clinical and laboratory data. *Eur J Intern Med* 2020; 76(6): 97-99.
9. Government of Pakistan. Official COVID-19 website. Available at <http://covid.gov.pk/> Accessed on 2020.
10. Chen T, Wu D, Chen H, Yan W, Yang D, Chen G, et al. Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. *Bio Med J* 2020; 368: m1091.
11. Lorente-Ros A, Monteagudo RJM, Rincón LM, Ortega PR, Rivas S, Martínez-Moya R, et al. Myocardial injury determination improves risk stratification and predicts mortality in COVID-19 patients. *Cardiol J* 2020; 10(1): 5603-10.
12. Wang L, He W, Yu X, Hu D, Bao M, Liu H, et al. Coronavirus Disease 2019 in elderly patients: characteristics and prognostic factors based on 4-week follow-up. *J Infect* 2020; 3(1): 19-25.
13. Wynants L, Van Calster B, Bonten MM, Collins GS, Debray TP, De Vos M, et al. Prediction models for diagnosis and prognosis of covid-19 infection: systematic review and critical appraisal. *Bio Med J* 2020; 1(1): 369-75.
14. Walter LA, McGregor AJ. Sex and gender specific observations and implications for COVID-19. *West J Emerg Med* 2020; 21(3): 507-9.
15. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020; 395(10229): 1054-62.
16. Hakeem R, Fawwad A. Diabetes in Pakistan: epidemiology, determinants and prevention. *J Diabetol* 2010; 1(3): 3-10.
17. Du RH, Liang LR, Yang CQ, Wang W, Cao TZ, Li M, et al. Predictors of mortality for patients with COVID-19 pneumonia caused by SARS-CoV-2: a prospective cohort study. *Eur Respir J* 2020; 55(1): 2000524.
18. Onder G, Rezza G, Brusaferro S. Case-Fatality rate and characteristics of patients dying in relation to COVID-19 in Italy. *J Am Med Assoc* 2020; 4683.
19. Ruan Q, Yang K, Wang W, Jiang L. Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Inten Care Med* 2020; 46(5): 846-48.