

## Platelet Indices as a Tool for Differentiation between Clonal Thrombocytosis and Reactive Thrombocytosis

Ammara Hafeez, Nasir ud Din Khattak, Samina Naeem\*, Helen Mary Robert, Shamaila Mohsin\*\*, Amal Mahmood

Department of Pathology, Combined Military Hospital, Lahore/National University of Medical Sciences (NUMS) Pakistan, \*Department of Pathology, Combined Military Hospital, Lahore Medical College/National University of Medical Sciences (NUMS) Pakistan, \*\*Department of Community Medicine, Army Medical College/National University of Medical Sciences (NUMS) Rawalpindi Pakistan

### ABSTRACT

**Objective:** To determine the efficacy of platelet indices as a tool for differentiation between clonal thrombocytosis and reactive thrombocytosis.

**Study Design:** Cross sectional study.

**Place and Duration of Study:** Department of Pathology, Combined Military Hospital, Lahore Pakistan, from Nov 2019 to Oct 2020.

**Methodology:** Fifty-six subjects were included in Reactive thrombocytosis group and 56 in Clonal thrombocytosis group. Fresh blood in EDTA anticoagulant was analyzed to determine complete blood counts and platelet parameters (Platelet distribution width, mean platelet volume, Platelet large cell ratio, Plateletcrit) using the automated hematological analyzer Sysmex KX-21.

**Results:** Mean age of the patients was 45.76 years with a range of 17-75 years. Assessing the cause of reactive thrombocytosis revealed that 20(35.7%) was infection and 16(28.5%) was iron deficiency anemia. In clonal thrombocytosis group 21(37.5%) were diagnosed with Chronic myeloid leukemia, 13(23.2%) with Essential thrombocythemia and 14(25%) with Polycythemia rubra vera. Platelet indices were compared in clonal thrombocytosis group and the reactive thrombocytosis group. Mean platelet volume, Platelet large cell ratio and Plateletcrit were significantly higher in the clonal thrombocytosis group ( $p=0.001$ ). Difference in Platelet distribution width was not found to be statistically significant in the two groups ( $p=0.07$ )

**Conclusion:** Platelet indices were higher in the clonal thrombocytosis group. Along with platelet count, they can serve as an efficient and cost-effective method in differentiating between clonal and reactive thrombocytosis.

**Keywords:** Anemia, Clonal thrombocytosis, Infection, Mean platelet volume, Plateletcrit, Platelet distribution width, Reactive thrombocytosis.

**How to Cite This Article:** Hafeez A, Khattak NUD, Naeem S, Robert HM, Mohsin S, Mahmood A. Platelet Indices as a Tool for Differentiation between Clonal Thrombocytosis and Reactive Thrombocytosis. *Pak Armed Forces Med J* 2023; 73(Suppl-1): S79-83. DOI: <https://doi.org/10.51253/pafmj.v73iSUPPL-1.6350>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## INTRODUCTION

Thrombocytosis is described as an increase in platelet count. Generally, a platelet count greater than  $450 \times 10^9/L$  is considered to be the working definition of thrombocytosis.<sup>1</sup> It usually comes across as an incidental laboratory finding when complete blood count is obtained for some unrelated reason. Thrombocytosis, when found creates an important diagnostic challenge for clinicians and pathologists alike.

The diagnostic workup for thrombocytosis starts with confirmation of presence of thrombocytosis by the evaluation of peripheral blood film. The next critical step in the diagnostic chain is to classify the condition into either primary/clonal or secondary/reactive thrombocytosis.<sup>2</sup> Various conditions contribute to the development of secondary (or reactive) thrombocytosis. These include iron deficiency anemia,

hemolysis, inflammation and malignancy.<sup>3</sup> Other causes include fever, infection, hemorrhage, post-operative thrombocytosis amongst other causes included in secondary thrombocytosis group. No age, gender, or race demographic predilection exists for secondary thrombocytosis. Secondary thrombocytosis is driven by overproduction of thrombopoietin, interleukin-6, other cytokines, or catecholamines in inflammatory, infectious, or neoplastic conditions or in situations of stress.<sup>4,5</sup>

Primary (clonal) thrombocytosis may be a result of various neoplasms of myeloproliferative origin. These can be further categorized into chronic myeloid leukemia (CML), essential thrombocythemia (ET), polycythemia rubra vera (PRV).<sup>6</sup> Gene mutations in Janus kinase 2 (JAK2), myeloproliferative leukemia virus oncogene (MPL) and/or Calreticulin (CALR) are usually found in a pathophysiological role in cases of primary thrombocytosis, leading to unchecked proliferative and anti-apoptotic effects on hematopoietic stem cells.

The morphology of platelets and their proliferation rate can be determined by automated

**Correspondence:** Dr Ammara Hafeez, Resident Hematology, Combined Military Hospital, Lahore, Pakistan  
Received: 15 Mar 2020; revision received: 10 Jun 2021; accepted: 21 Jun 2021

complete blood count analysers. Platelet indices include plateletcrit (PCT), mean platelet volume (MPV), and platelet distribution width (PDW).<sup>7</sup> along with Platelet Large cell ratio (P-LCR). Variations in the platelet size, morphology and volume are determined by PDW. It is usually raised in case of platelet anisocytosis.<sup>8</sup> PCT is the volume occupied by platelets as a percentage and is calculated according to the formula  $PCT = \text{platelet count} \times \text{MPV} / 10,000$ . Mean platelet volume (MPV) is a measurement of average size of platelet cells in blood.<sup>9</sup>

Given the cost effectiveness of determination of platelet indices as compared to genetic testing, the aim of this study is to evaluate the effectiveness of platelet indices as a diagnostic tool in order to differentiate between clonal and reactive thrombocytosis.

## METHODOLOGY

The cross-sectional study was carried out at the Department of Pathology Combined Military Hospital Lahore, Pakistan over a timeframe of one year from November 2019 to October 2020. Permission from the ethical review board (Ltr no. 159/2020) was taken prior to conducting the study. Informed consent was taken from each patient. A sample size of 112 was calculated using OpenEpi sample size calculator taking mean MPV  $9.8 \pm 2.0$  fL in clonal thrombocytosis and  $11.0 \pm 2.8$  fL in reactive thrombocytosis group.<sup>10</sup>

**Inclusion Criteria:** Subjects of either gender, aged  $\geq 12$  years having platelet count  $>400 \times 10^9/L$  were included.

**Exclusion Criteria:** Patients receiving growth factors such as thrombopoietin receptor agonists were excluded.

Fifty-six (56) patients were included in Reactive thrombocytosis group and 56 patients in Clonal thrombocytosis group based on history (fever, anemia, blood loss, thrombotic episodes, myocardial infarction, CVA, erythromyalgia), relevant physical examination (splenomegaly, hepatomegaly, lymphadenopathy) and laboratory investigations.

Patients fulfilling WHO criteria for Myeloproliferative neoplasms (including Chronic Myeloid Leukemia, Polycythemia Rubra Vera, Essential Thrombocythemia, Primary Myelofibrosis etc) were labeled as clonal thrombocytosis based primarily on findings of bone marrow examination, genetic studies and serum erythropoietin levels.<sup>11</sup> Patients with history of fever, signs of localized or systemic infection, iron deficiency anemia, hemorrhage, chronic inflammatory

diseases, malignancy, and history of surgery were included in reactive thrombocytosis group.

Five (05) milliliters of fresh blood in a tube containing EDTA anticoagulant was analyzed to determine complete blood counts (Total Leucocyte count (TLC), Hemoglobin (Hb), Mean corpuscular volume (MCV), Mean corpuscular hemoglobin (MCH), Mean corpuscular hemoglobin concentration (MCHC), Platelet count, Differential leucocyte count (DLC) including neutrophils, lymphocytes, monocytes, eosinophils, basophils, myelocytes, metamyelocytes, blast cells) using automated hematological analyser Sysmex KX-21. Platelet parameters including Platelet Distribution Width (PDW), Mean Platelet Volume (MPV), Platelet Large cell Ratio (P-LCR), Plateletcrit (PCT) were analysed using same hematological analyser which uses impedance counting as the principle for hematological analysis.

Statistical analysis was done using SPSS version 26.0. Mean and SD were calculated for quantitative variables and frequency and percentages for qualitative variables. Mean values of the complete blood picture and platelet indices were compared using independent samples t test between the two groups. Difference in the clinical parameters were analysed using Chi square test between the two groups. The  $p$  value of  $\leq 0.05$  was considered to be statistically significant. ROC (receiver operating characteristic) curve was used to assess the cut off value for sensitivity and specificity of platelet indices in differentiating reactive and clonal thrombocytosis.


## RESULTS

Out of a total of 112 patients, 42(37.5%) were females and 70(62.5%) were males. Mean age of the patients was 45.76 years with a standard deviation of 16.74 years and a range of 17-75 years. Assessing the cause of reactive thrombocytosis revealed that in 20 (35.7%) cases it was infection and in 16(28.5%) it was iron deficiency anemia. In clonal thrombocytosis group, 21(37.5%) were diagnosed with chronic myeloid leukemia (CML), 13(23.2%) with essential thrombocythemia (ET) and 14(25%) with polycythemia rubra vera (PRV). There was a statistically significant difference in the mean value of TLC, Hb and platelet count between the two groups.

On analyzing differential leukocyte count, there was a statistically increase in monocytes, basophils and myelocytes in the clonal thrombocytosis group as compared to the reactive thrombocytosis group ( $p < 0.05$ ).


**Table: Mean value of blood cells in the Reactive Thrombocytosis and Clonal Thrombocytosis Group (n=112)**

Parameter	Reactive Thrombocytosis n=56	Clonal Thrombocytosis n=56	p-value
Total leucocyte count×10 <sup>9</sup> /L	11.68±5.0	86.2±112.7	0.001
Hemoglobin g/dL	10.39±1.9	12.7±3.8	0.001
Mean Corpuscular Volume fL	82.17±7.7	82.04±11.0	0.03
Mean Corpuscular Hemoglobin pg	26.84±3.2	27.10±5.5	0.005
Mean Corpuscular Hemoglobin Concentration g/dL	32.57±1.5	32.62±3.37	0.001
Platelet count×10 <sup>9</sup> /L	569.13±120.3	821.95±513.2	0.001



**Figure-1: Differential Leukocyte count in Clonal and Reactive Thrombocytosis Groups (n=112)**

Platelet indices were compared in clonal thrombocytosis group and the reactive thrombocytosis group. MPV, P-LCR and PCT were significantly higher in the clonal thrombocytosis group ( $p=0.001$ ). The difference in PDW was not statistically significant in both the groups ( $p=0.07$ ).



Area under the curve for PCT was 0.372, PDW was 0.41 and MPV was 0.50.

**Figure-2: ROC Curve showing Specificity And Sensitivity of MPV, PCT and PDW in clonal and Reactive Thrombocytosis (n=112)**

Area under the curve for PCT was 0.372, PDW was 0.41 and MPV was 0.50.

## DISCUSSION

Platelets are derived from bone marrow precursor cells called megakaryocytes. They have a diameter of 3-5  $\mu\text{m}$  and a volume of 4.5-11 fL. Protoplatelets are inactivated forms of platelets in the bloodstream. These cells are anucleate and discoidal in shape. Platelet activation results in change in the morphology from discoid to dendritic shape. Prothrombotic molecules released from alpha and dense granules lead to aggregation and further recruitment of platelets.

Thrombocytosis is an increase in platelet count from its normal range (150-450×10<sup>9</sup>/L. A platelet count of >450×10<sup>9</sup>/L qualifies as thrombocytosis.

Establishing the differentiation between clonal and reactive causes of thrombocytosis has significant therapeutic implications. Clonal thrombocytosis has been linked with increased incidence of thromboembolic and bleeding tendency and complications according to a study by Andrew I. Schafer,<sup>4</sup> whereas secondary thrombocytosis usually does not show any such association. Cerebral, coronary and hepatic vessels are mostly involved. Thus, prophylaxis for prevention of thromboembolic events in high-risk patients falling in clonal thrombocytosis group is of utmost importance.

The most common cause of reactive thrombocytosis in our study population was infection (35.7%) followed by iron deficiency anemia (28.5%). A study by Khalid Nafih and co-workers showed infection (27%) as the leading cause of thrombocytosis, followed by malignancies of non hematological type, iron deficiency anemia, hemolysis, postoperative thrombocytosis and vascular defects to name a few.<sup>12</sup> According to Syed Naveen and colleagues, causes of thrombocytosis included infections (44.9%), rebound thrombocytosis, tissue injury, myeloproliferative neoplasms, iron deficiency anemia, malignancy, post splenectomy, etc.<sup>13</sup>

Previous studies on the subject showed male predominance in the frequency of thrombocytosis in adults as well as in children.<sup>14,15</sup> This correlates with our study as well showing 62.5% males and 37.5% females from a total sample of 112 subjects.

Findings of our study showed an increased platelet distribution width (PDW), mean platelet volume (MPV), plateletcrit (PCT) and platelet large cell ratio (P-LCR) in the clonal thrombocytosis group as compared to the reactive thrombocytosis group. Our study showed the mean PDW value was 12.14±2.6 fL in the

clonal thrombocytosis group and  $11.25 \pm 1.6$  in the reactive thrombocytosis group which was a statistically insignificant difference ( $p=0.07$ ). AUC for PCT was 0.372, PDW was 0.41 and MPV was 0.50. Tanzeel and colleagues revealed that mean PDW was significantly higher in the clonal thrombocytosis group as compared to reactive thrombocytosis group and the AUC was significantly higher for MPV (0.989) and PDW (0.981) as compared to the other study.<sup>16</sup> A study done by Sangeetha *et al.* showed a significantly raised PDW value in thrombocytosis group as compared to the normal control group. Rachita *et al.* revealed contradictory results and suggested that with the increasing number of platelets there was a decreasing trend in MPV, P-LCR and PDW.<sup>17</sup> Similar results were portrayed by Sabramaniam in his study increasing platelet counts were associated with decrease in MPV ( $p=0.001$ ).<sup>14</sup>

A study conducted by Tafazzoli *et al.* concluded that platelet count of patients with primary thrombocytosis was greater than  $1000 \times 10^9/L$  as compared to secondary thrombocytosis and a higher mean platelet volume (MPV) was seen in primary thrombocytosis group ( $p=0.007$ ).<sup>18</sup> In our study the platelet count was  $569.13 \pm 120.3 \times 10^9/L$  in the reactive thrombocytosis group and  $821.95 \pm 513.2$  in the clonal thrombocytosis group. In a study carried out by Khaleel *et al.* in Iraq, findings in accordance with our study were revealed. They concluded that MPV, PCT and PDW were appreciably raised in the primary thrombocytosis group as compared to the secondary thrombocytosis group.<sup>19</sup>

In our study the value of PDW was not significantly different between the two groups. This was in contrast to the study conducted by Golajapu *et al.* who concluded that MPV, PDW and PCT values were significantly higher in the clonal group as compared to the reactive thrombocytosis group.<sup>20</sup> Nafih and co-workers also suggested that MPV and PDW both were raised in the reactive thrombocytosis group.<sup>12</sup> Increase in the MPV may lead to a series of thrombotic complications which are more commonly linked to primary thrombocytosis group.

The reason behind this is that the large platelets are more enzymatically and metabolically active, leading to greater thrombotic activity.<sup>21</sup> MPV and PDW both have a direct relationship and they usually change in the same direction.<sup>22</sup> The diagnostic role of PDW in differentiating clonal and reactive thrombocytosis group is some what contradictory and was not found to be of any significance in our study.

## CONCLUSION

Platelet indices; MPV, PCT and P-LCR are higher in the primary (or clonal) thrombocytosis group. Along with platelet counts, they can serve as an efficient and cost-effective method in differentiating clonal and reactive thrombocytosis group.

Secondary Thrombocytosis occurs mostly as a result of infection or inflammation, anemia and hemorrhage along with certain other causes. Clonal (or primary) thrombocytosis is generally part of a myeloproliferative process and is usually confirmed with bone marrow examination and genetic testing. This study highlights the importance of platelet indices as a sensitive predictive marker to distinguish between clonal and reactive thrombocytosis, thereby reducing the need or urgency of invasive or expensive procedures.

**Conflict Of Interest:** None.

## Author's Contribution

Following authors have made substantial contributions to the manuscript as under:

AH: & NUDK: Study design, drafting the manuscript, data interpretation, critical review, approval of the final version to be published.

SN: & HMR: Critical review, concept, drafting the manuscript, approval of the final version to be published.

SM: & AM: Data acquisition, data analysis, approval of the final version to be published.

Authors agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

## REFERENCES

1. Vo QT, Thompson DF. A review and assessment of drug-induced thrombocytosis. *Ann Pharmacother.* 2019; 53(5): 523-536. <https://doi.org/10.1177/1060028018819450>
2. Thakral B, Wang SA. Thrombocytosis. In: Wang SA, Hasserjian R. (eds) *Diagnosis of Blood and Bone Marrow Disorders.* New York, United States of America. Spring Cham. 2018; 1(01): 225-242.
3. Klose N, Boyle S, Harvey Y, Niemann J. Aetiology and differentiating characteristics of primary versus secondary thrombocytosis the clot thickens. *Intern Med J* 2019; 51(1): 124
4. Schafer AL. Thrombocytosis. *N Engl J Med* 2004; 350(12): 1211-1219.
5. Araneda M, Krishnan V, Hall K, Kalbfleisch J, Krishnaswamy G, Krishnan K. Reactive and clonal thrombocytosis: proinflammatory and hematopoietic cytokines and acute phase proteins. *South Med J* 2001; 94(4): 417-420.
6. Hsieh RW, Ravindran A, Hook CC, Begna KH, Ashrani AA, Pruthi RK, et al. Etiologies of extreme thrombocytosis: a contemporary series. *Mayo Clin. Proc* 2019; 94(8): 1542-1550.
7. Pogorzelska K, Krętońska A, Krawczuk-Rybak M, Sawicka-Żukowska M. Characteristics of platelet indices and their prognostic significance in selected medical condition—a systematic review. *Adv Med Sci* 2020; 65(2): 310-5. <https://doi.org/10.1016/j.advms.2020.05.002>
8. Budak YU, Polat M, Huysal K. The use of platelet indices, plateletcrit, mean platelet volume and platelet distribution width in emergency non-traumatic abdominal surgery: a systematic review. *Biochem Med* 2016; 26(2): 178-193.



## Clonal Thrombocytosis and Reactive Thrombocytosis

9. Beyan C, Kaptan K, Ifran A. Platelet count, mean platelet volume, platelet distribution width, and plateletcrit do not correlate with optical platelet aggregation responses in healthy volunteers. *J thromb thrombolysis* 2006; 22(3): 161-164
  10. Kim MJ, Park PW, Seo YH, Kim KH, Seo JY, Jeong JH, et. al. Comparison of platelet parameters in thrombocytopenic patients associated with acute myeloid leukemia and primary immune thrombocytopenia. *Blood Coagul Fibrinol* 2014; 25(3): 221-225.
  11. Swerdlow SH, Campo E, Harris NL, Jaffe ES, Pileri SA, Stein H, et. al. WHO Classification of Tumours of Haematopoietic and Lymphoid Tissues, 4th edition; IARC: Lyon, France, 2017, [internet] available at : <https://publications.iarc.fr/Book-And-Report-Series/Who-Classification-Of-Tumours/WHO-cation-Of-Tumours-Of-Haematopoietic-And-Lymphoid-Tissues-2017>
  12. Nafih K, Taib SM, Saeed BA. Platelet indices in the differential diagnosis of thrombocytosis. *Ann Coll Med* 2009; 35(1): 33-36
  13. Syed NN, Usman M, Khurshid M. Thrombocytosis: age dependent aetiology and analysis of platelet indices for differential diagnosis. *Indian J Pathol Microbiol* 2007; 50(3): 628-633.
  14. Subramaniam N, Mundkur S, Kini P, Bhaskaranand N, Aroor S. Clinicohematological study of thrombocytosis in children. *ISRN Hematol*. 2014; 2014: 389257. <https://doi: 10.1155/2014/389257>.
  15. Lee E, Kim M, Jeon K, Lee J, Lee JS, Kim HS et. al. Mean platelet volume, platelet distribution width, and platelet count, in connection with immune thrombocytopenic purpura and essential thrombocytopenia. *Lab Med* 2019; 50(3): 279-285.
  16. Imran T, Zafar L, Altaf HN, Orakzai S, Butt T, Bakhtiar F. Evaluation of platelet parameters (PDW and MPV) for differentiation between clonal thrombocytosis and reactive thrombocytosis. *Rawal Med J* 2017; 42(3): 302-305
  17. Sarangi R, Pradhan S, Dhanawat A, Patanayak R, Benia G. Thrombocytosis in children: Clinico-hematological profile from a single centre in Eastern India *J Lab Physicians* 2018; 10(1): 34-37.
  18. Tafazzoli M, Keramati MR, Vakili R. Etiology of thrombocytosis and the use of platelet parameters to distinguish between clonal and reactive thrombocytosis. *Int J Hematol Oncol* 2006; 29(4): 071-076
  19. Khaleel KJ, Ahmed AA, Alwash MM, Yasin N. Mean platelet volume, platelet distribution width and plateletcrit values in differentiating clonal from secondary thrombocytosis. *Med J Babylon* 2013; 10(2): 473-478.
  20. Golajapu R, Paul T, Uppin MS, Shantveer GU, Narendra AS, Radhika, et al. Utility of Platelet Indices to Differentiate Reactive vs Clonal Thrombocytosis. *Int J Sci Res (Raipur)* 2019; 8(6): 1729-1733.
  21. Bolayır A, Gökçe ŞF. The role of mean platelet volume, platelet distribution width and platelet/lymphocyte ratio in development of cerebral venous thrombosis. *Cumhuriyet Tıp Dergisi* 2017; 39(4): 683-691.
  22. Subramanian M, Muthuvel E, Srinivasan C, Sekar G. Thrombocytopenia Vs Pseudo thrombocytopenia Will indices help us???. *Int J Res Pharm Sci* 2020; 11(SPL2): 108-112.
-